

What is Claimed:

- 1. An isolated polynucleotide comprising a sequence selected from the group consisting of:
- (a) sequences provided in SEQ ID NO: 1-111, 115-171, 173-175, 177, 179-305, 307-315, 326, 328, 330, 332-335, 340-375, 381, 382 and 384-476, 524, 526, 530, 531, 533, 535, 536, 552, 569-572, 587, 591, 593-606, 618-705, 709-774, 777, 789, 817, 823, 824, 878, 880-882, 894, 896, 907, 908, 916-919 and 929-931;
- (b) complements of the sequences provided in SEQ ID NO: 1-111, 115-171, 173-175, 177, 179-305, 307-315, 326, 328, 330, 332-335, 340-375, 381, 382 and 384-476, 524, 526, 530, 531, 533, 535, 536, 552, 569-572, 587, 591, 593-606, 618-705, 709-774, 777, 789, 817, 823, 824, 878, 880-882, 894, 896, 907, 908, 916-919 and 929-931;
- (c) sequences consisting of at least 20 contiguous residues of a sequence provided in SEQ ID NO: 1-111, 115-171, 173-175, 177, 179-305, 307-315, 326, 328, 330, 332-335, 340-375, 381, 382 and 384-476, 524, 526, 530, 531, 533, 535, 536, 552, 569-572, 587, 591, 593-606, 618-705, 709-774, 777, 789, 817, 823, 824, 878, 880-882, 894, 896, 907, 908, 916-919 and 929-931;
- (d) sequences that hybridize to a sequence provided in SEQ ID NO: 1-111, 115-171, 173-175, 177, 179-305, 307-315, 326, 328, 330, 332-335, 340-375, 381, 382 and 384-476, 524, 526, 530, 531, 533, 535, 536, 552, 569-572, 587, 591, 593-606, 618-705, 709-774, 777, 789, 817, 823, 824, 878, 880-882, 894, 896, 907, 908, 916-919 and 929-931 under moderately stringent conditions;
- (e) sequences having at least 75% identity to a sequence of SEQ ID NO: 1-111, 115-171, 173-175, 177, 179-305, 307-315, 326, 328, 330, 332-335, 340-375, 381, 382 and 384-476, 524, 526, 530, 531, 533, 535, 536, 552, 569-572, 587, 591, 593-606,





618-705, 709-774, 777, 789, 817, 823, 824, 878, 880-882, 894, 896, 907, 908, 916-919 and 929-931;

- (f) sequences having at least 90% identity to a sequence of SEQ ID NO: 1-111, 115-171, 173-175, 177, 179-305, 307-315, 326, 328, 330, 332-335, 340-375, 381, 382 and 384-476, 524, 526, 530, 531, 533, 535, 536, 552, 569-572, 587, 591, 593-606, 618-705, 709-774, 777, 789, 817, 823, 824, 878, 880-882, 894, 896, 907, 908, 916-919 and 929-931; and
- (g) degenerate variants of a sequence provided in SEQ ID NO: 1-111, 115-171, 173-175, 177, 179-305, 307-315, 326, 328, 330, 332-335, 340-375, 381, 382 and 384-476, 524, 526, 530, 531, 533, 535, 536, 552, 569-572, 587, 591, 593-606, 618-705, 709-774, 777, 789, 817, 823, 824, 878, 880-882, 894, 896, 907, 908, 916-919 and 929-931.

An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- (a) sequences recited in SEQ ID NO: 112-114, 172, 176, 178, 327, 329, 331, 336, 339, 376-380, 383, 477-483, 496, 504, 505, 519, 520, 522, 525, 527, 532, 534, 537-551, 553-568, 573-586, 588-590, 592, 706-708, 775, 776, 778, 780, 781, 811, 814, 818, 826, 827, 853, 855, 858, 860-862, 866-877, 879, 883-893, 895, 897, 898, 909-915, 920-928 and 932-934;
- (b) sequences having at least 70% identity to a sequence of SEQ ID NO: 112-114, 172, 176, 178, 327, 329, 331, 336, 339, 376-380, 383, 477-483, 496, 504, 505, 519, 520, 522, 525, 527, 532, 534, 537-551, 553-568, 573-586, 588-590, 592, 706-708, 775, 776, 778, 780, 781, 811, 814, 818, 826, 827, 853, 855, 858, 860-862, 866-877, 879, 883-893, 895, 897, 898, 909-915, 920-928 and 932-934;
- (c) sequences having at least 90% identity to a sequence of SEQ ID NO: 112-114, 172, 176, 178, 327, 329, 331, 336, 339, 376-380, 383, 477-483, 496, 504, 505, 519, 520, 522, 525, 527, 532, 534, 537-551, 553-568, 573-586, 588-590, 592, 706-708, 775, 776, 778, 780, 781, 811, 814, 818, 826, 827, 853, 855, 858, 860-862, 866-877, 879, 883-893, 895, 897, 898, 909-915, 920-928 and 932-934;





- (d) sequences encoded by a polynucleotide of claim 1;
- (e) sequences having at least 70% identity to a sequence encoded by a polynucleotide of claim 1; and
- (f) sequences having at least 90% identity to a sequence encoded by a polynucleotide of claim 1.
- 3. An expression vector comprising a polynucleotide of claim 1 operably linked to an expression control sequence.

4. A host cell transformed or transfected with an expression vector according to claim 3.

- 5. An isolated antibody, or antigen-binding fragment thereof, that specifically binds to a polypeptide of claim 2.
- 6. A method for detecting the presence of a cancer in a patient, comprising the steps of:
 - (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with a binding agent that binds to a polypeptide of claim 2;
- (c) detecting in the sample an amount of polypeptide that binds to the binding agent; and
- (d) comparing the amount of polypeptide to a predetermined cut-off value and therefrom determining the presence of a cancer in the patient.
- 7. A fusion protein comprising at least one polypeptide according to claim 2.

claim 2.





- 8. An oligonucleotide that hybridizes to a sequence recited in SEQ ID NO: 1-111, 115-171, 173-175, 177, 179-305, 307-315, 326, 328, 330, 332-335, 340-375, 381, 382 and 384-476, 524, 526, 530, 531, 533, 535, 536, 552, 569-572, 587, 591, 593-606, 618-705, 709-774, 777, 789, 817, 823, 824, 878, 880-882, 894, 896, 907, 908, 916-919 and 929-931 under moderately stringent conditions.
- 9. A method for stimulating and/or expanding T cells specific for a tumor protein, comprising contacting T cells with at least one component selected from the group consisting of:
 - (a) polypeptides according to claim 2;
 - (b) polynucleotides according to claim 1; and
- (c) antigen-presenting cells that express a polypeptide according to claim 1,

under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells.

- 10. An isolated T cell population, comprising T cells prepared according to the method of claim 9.
- 11. A composition comprising a first component selected from the group consisting of physiologically acceptable carriers and immunostimulants, and a second component selected from the group consisting of:
 - (a) polypeptides according to claim 2;
 - (b) polynucleotides according to claim 1;
 - (c) antibodies according to claim 5;
 - (d) fusion proteins according to claim 7;
 - (e) T cell populations according to claim 10; and
 - (f) antigen presenting cells that express a polypeptide according to





- 12. A method for stimulating an immune response in a patient, comprising administering to the patient a composition of claim 11.
- 13. A method for the treatment of a cancer in a patient, comprising administering to the patient a composition of claim 11.

A method for determining the presence of a cancer in a patient, comprising the steps of:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide according to claim 8;
- (c) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; and
- (d) compare the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefrom determining the presence of the cancer in the patient.
- 15. A diagnostic kit comprising at least one oligonucleotide according to claim 8.
- 16. A diagnostic kit comprising at least one antibody according to claim 5 and a detection reagent, wherein the detection reagent comprises a reporter group.
- 17. A method for inhibiting the development of a cancer in a patient, comprising the steps of:
- (a) incubating CD4+ and/or CD8+ T cells isolated from a patient with at least one component selected from the group consisting of: (i) polypeptides according to claim 2; (ii) polynucleotides according to claim 1; and (iii) antigen presenting cells that express a polypeptide of claim 2, such that T cell proliferate;





(b) administering to the patient an effective amount of the proliferated T cells,

and thereby inhibiting the development of a cancer in the patient.

ADD AG)